

# 鄰苯二甲酸酐 (苯酐) PA Phthalic Anhydride

### 說明 Description

鄰苯二甲酸酐是由鄰二甲苯經部份氧化反應而 成。在室溫為白色結晶片狀形態,與水接觸會形 成酸。鄰苯二甲酸酐主要用以製造鄰苯二甲酸酯 類可塑劑,供作為PVC加工之用。

Phthalic anhydride (PA) is produced via partial oxidation of orth-xylene. At room temperature, it forms white crystal-like flakes. When contacted with water, phthalic acid is produced. Phthalic anhydride is widely used to make phthalate esters plasticizers for PVC processing.

CAS No.: 85-44-9

規格 Specifications			Test Method
凝 固 點 Solidification Point	°C	≧ 130.8	ASTM-D-1493
純 度 Purity	wt %	≧ 99.8	GC
酜酸含量 Phthalic Acid	wt %	< 0.50	CNS-12231
馬 來 酐 Maleic Anhydride	wt %	< 0.05	GC
熔融色相 Molten Color	APHA	≦ 20	ASTM-D-3366
加熱安定性 Heat Hazen Number	APHA	≦ 50	ASTM-D-3366

### 物性 Physical Properties

分子式 Molecular Formula

構造式 Structural Formula

C<sub>8</sub>H<sub>4</sub>O<sub>3</sub>

g/mole 148.12 分子量 Molecular Weight 沸 點 Boiling Point (760mmHg) °C 285 閃火點 Flash Point °C 152

### 包裝 Packing

液態槽車;

袋裝:750kg、600kg、500kg、25kg

Molten: Tank Truck;

Flake Bag: 750kg \ 600kg \ 500kg \ 25kg

(with Pallets)

## 用途 Usage

- 可塑劑原料
- 不飽和聚酯樹酯原料、醇酸樹酯原料
- •染料、顏料、驅蟲劑、耐燃劑、…等原料
- Raw material of phthalate plasticizers
- Unsaturated polyester and Alkyd resin.
- Dyes > Pigments > Insecticides > Fire retardants > · · · etc

#### 聯絡人 Contact:

賴宏昇 Horng-Sheng Lai

+886 2 2712 2211 EXT.5721 eapublic@npc.com.tw

林訓民 Fred Lin

fredlin@npc.com.tw

+886 2 2712 2211 EXT.5723

#### 南亞塑膠工業股份有限公司 NAN YA PLASTICS CORPORATION

化工第一事業部・PETROCHEMICAL 1ST. DIV.

台北市敦化北路201號 電話:(02)2712-2211 EXT 8127 201, TUN HWA N. ROAD, TAIPEI, TAIWAN, R.O.C. TEL: (886-2) 2712-2211 EXT 8127 FAX: (886-2) 2717-8534

14